



We claim:

1. A process for preparing 2-cyclopentenones of the general  
5 formula:



- 15 where R<sup>1</sup> to R<sup>4</sup> are each hydrogen atoms or are alkyl or alkenyl radicals having from 1 to 12 carbon atoms, cycloalkyl or cycloalkenyl radicals having from 5 to 7 carbon atoms, aralkylene or aryl radicals, by converting hexenedioic acids and/or their esters of the general formulae



- 25 where R<sup>1</sup> to R<sup>4</sup> are each as defined above and R<sup>5</sup> and R<sup>6</sup> are each hydrogen atoms or are alkyl radicals having from 1 to 12 carbon atoms, cycloalkyl radicals having 5 or 6 carbon atoms, aralkyl or aryl radicals, at temperatures of from 150 to  
30 450°C, over solid, oxidic catalysts, wherein the catalysts on an oxidic support material comprise from 0.01 to 5% by weight of at least one alkali metal oxide.

- 35 2. A process as claimed in claim 1, wherein the support material used is a metal oxide of main groups II to V, transition groups I to VIII of the Periodic Table of the Elements, an oxide of the rare earth metals or a mixture thereof.
- 40 3. A process as claimed in claim 1, wherein the support material used is aluminum oxide and/or silicon oxide.
- 45 4. A process as claimed in claim 1, wherein sodium oxide and/or potassium oxide are used.

9

5. A process as claimed in claim 1, wherein the reaction is carried out in a fixed bed.

6. A process as claimed in claim 1, wherein the starting materials of the general formulae II and III are prepared by dimerizing acrylic esters.

10

15

20

25

30

35

40

45